

Using Properties and Known Values to Find Other Integrals

9. Suppose that f and g are integrable and that

$$\int_1^2 f(x) dx = -4, \int_1^5 f(x) dx = 6, \int_1^5 g(x) dx = 8.$$

Use the rules in Table 5.3 to find

a. $\int_2^2 g(x) dx$	b. $\int_5^1 g(x) dx$
c. $\int_1^2 3f(x) dx$	d. $\int_2^5 f(x) dx$
e. $\int_1^5 [f(x) - g(x)] dx$	f. $\int_1^5 [4f(x) - g(x)] dx$

10. Suppose that f and h are integrable and that

$$\int_1^9 f(x) dx = -1, \int_7^9 f(x) dx = 5, \int_7^9 h(x) dx = 4.$$

Use the rules in Table 5.3 to find

a. $\int_1^9 -2f(x) dx$	b. $\int_7^9 [f(x) + h(x)] dx$
c. $\int_7^9 [2f(x) - 3h(x)] dx$	d. $\int_9^1 f(x) dx$
e. $\int_1^7 f(x) dx$	f. $\int_9^7 [h(x) - f(x)] dx$

11. Suppose that $\int_1^2 f(x) dx = 5$. Find

a. $\int_1^2 f(u) du$	b. $\int_1^2 \sqrt{3}f(z) dz$
c. $\int_2^1 f(t) dt$	d. $\int_1^2 [-f(x)] dx$

12. Suppose that $\int_{-3}^0 g(t) dt = \sqrt{2}$. Find

a. $\int_0^{-3} g(t) dt$	b. $\int_{-3}^0 g(u) du$
c. $\int_{-3}^0 [-g(x)] dx$	d. $\int_{-3}^0 \frac{g(r)}{\sqrt{2}} dr$

13. Suppose that f is integrable and that $\int_0^3 f(z) dz = 3$ and $\int_0^4 f(z) dz = 7$. Find

a. $\int_3^4 f(z) dz$	b. $\int_4^3 f(t) dt$
-----------------------	-----------------------

14. Suppose that h is integrable and that $\int_{-1}^1 h(r) dr = 0$ and $\int_{-1}^3 h(r) dr = 6$. Find

a. $\int_1^3 h(r) dr$	b. $-\int_3^1 h(u) du$
-----------------------	------------------------

Using Area to Evaluate Definite Integrals

In Exercises 15–22, graph the integrands and use areas to evaluate the integrals.

15. $\int_{-2}^4 \left(\frac{x}{2} + 3\right) dx$	16. $\int_{1/2}^{3/2} (-2x + 4) dx$
---	-------------------------------------

17. $\int_{-3}^3 \sqrt{9 - x^2} dx$

19. $\int_{-2}^1 |x| dx$

21. $\int_{-1}^1 (2 - |x|) dx$

18. $\int_{-4}^0 \sqrt{16 - x^2} dx$

20. $\int_{-1}^1 (1 - |x|) dx$

22. $\int_{-1}^1 (1 + \sqrt{1 - x^2}) dx$

Use areas to evaluate the integrals in Exercises 23–26.

23. $\int_0^b \frac{x}{2} dx, b > 0$

24. $\int_0^b 4x dx, b > 0$

25. $\int_a^b 2s ds, 0 < a < b$

26. $\int_a^b 3t dt, 0 < a < b$

Evaluations

Use the results of Equations (1) and (3) to evaluate the integrals in Exercises 27–38.

27. $\int_1^{\sqrt{2}} x dx$

28. $\int_{0.5}^{2.5} x dx$

29. $\int_{\pi}^{2\pi} \theta d\theta$

30. $\int_{\sqrt{2}}^{5\sqrt{2}} r dr$

31. $\int_0^{\sqrt[3]{7}} x^2 dx$

32. $\int_0^{0.3} s^2 ds$

33. $\int_0^{1/2} t^2 dt$

34. $\int_0^{\pi/2} \theta^2 d\theta$

35. $\int_a^{2a} x dx$

36. $\int_a^{\sqrt{3}a} x dx$

37. $\int_0^{\sqrt[3]{b}} x^2 dx$

38. $\int_0^{3b} x^2 dx$

Use the rules in Table 5.3 and Equations (1)–(3) to evaluate the integrals in Exercises 39–50.

39. $\int_3^1 7 dx$

40. $\int_0^{-2} \sqrt{2} dx$

41. $\int_0^2 5x dx$

42. $\int_3^5 \frac{x}{8} dx$

43. $\int_0^2 (2t - 3) dt$

44. $\int_0^{\sqrt{2}} (t - \sqrt{2}) dt$

45. $\int_2^1 \left(1 + \frac{z}{2}\right) dz$

46. $\int_3^0 (2z - 3) dz$

47. $\int_1^2 3u^2 du$

48. $\int_{1/2}^1 24u^2 du$

49. $\int_0^2 (3x^2 + x - 5) dx$

50. $\int_1^0 (3x^2 + x - 5) dx$

Finding Area

In Exercises 51–54 use a definite integral to find the area of the region between the given curve and the x -axis on the interval $[0, b]$.

51. $y = 3x^2$

52. $y = \pi x^2$

53. $y = 2x$

54. $y = \frac{x}{2} + 1$

Average Value

In Exercises 55–62, graph the function and find its average value over the given interval.

55. $f(x) = x^2 - 1$ on $[0, \sqrt{3}]$

56. $f(x) = -\frac{x^2}{2}$ on $[0, 3]$ 57. $f(x) = -3x^2 - 1$ on $[0, 1]$

58. $f(x) = 3x^2 - 3$ on $[0, 1]$

59. $f(t) = (t - 1)^2$ on $[0, 3]$

60. $f(t) = t^2 - t$ on $[-2, 1]$

61. $g(x) = |x| - 1$ on a. $[-1, 1]$, b. $[1, 3]$, and c. $[-1, 3]$

62. $h(x) = -|x|$ on a. $[-1, 0]$, b. $[0, 1]$, and c. $[-1, 1]$

Theory and Examples

63. What values of a and b maximize the value of

$$\int_a^b (x - x^2) dx$$

(Hint: Where is the integrand positive?)

64. What values of a and b minimize the value of

$$\int_a^b (x^4 - 2x^2) dx$$

65. Use the Max-Min Inequality to find upper and lower bounds for the value of

$$\int_0^1 \frac{1}{1+x^2} dx.$$

66. (Continuation of Exercise 65) Use the Max-Min Inequality to find upper and lower bounds for

$$\int_0^{0.5} \frac{1}{1+x^2} dx \quad \text{and} \quad \int_{0.5}^1 \frac{1}{1+x^2} dx.$$

Add these to arrive at an improved estimate of

$$\int_0^1 \frac{1}{1+x^2} dx.$$

67. Show that the value of $\int_0^1 \sin(x^2) dx$ cannot possibly be 2.

68. Show that the value of $\int_1^0 \sqrt{x+8} dx$ lies between $2\sqrt{2} \approx 2.8$ and 3.

69. **Integrals of nonnegative functions** Use the Max-Min Inequality to show that if f is integrable then

$$f(x) \geq 0 \quad \text{on} \quad [a, b] \quad \Rightarrow \quad \int_a^b f(x) dx \geq 0.$$

70. **Integrals of nonpositive functions** Show that if f is integrable then

$$f(x) \leq 0 \quad \text{on} \quad [a, b] \quad \Rightarrow \quad \int_a^b f(x) dx \leq 0.$$

71. Use the inequality $\sin x \leq x$, which holds for $x \geq 0$, to find an upper bound for the value of $\int_0^1 \sin x dx$.

72. The inequality $\sec x \geq 1 + (x^2/2)$ holds on $(-\pi/2, \pi/2)$. Use it to find a lower bound for the value of $\int_0^1 \sec x dx$.

73. If $\text{av}(f)$ really is a typical value of the integrable function $f(x)$ on $[a, b]$, then the number $\text{av}(f)$ should have the same integral over $[a, b]$ that f does. Does it? That is, does

$$\int_a^b \text{av}(f) dx = \int_a^b f(x) dx?$$

Give reasons for your answer.

74. It would be nice if average values of integrable functions obeyed the following rules on an interval $[a, b]$.

a. $\text{av}(f + g) = \text{av}(f) + \text{av}(g)$

b. $\text{av}(kf) = k \text{av}(f)$ (any number k)

c. $\text{av}(f) \leq \text{av}(g)$ if $f(x) \leq g(x)$ on $[a, b]$.

Do these rules ever hold? Give reasons for your answers.

75. Use limits of Riemann sums as in Example 4a to establish Equation (2).

76. Use limits of Riemann sums as in Example 4a to establish Equation (3).

77. Upper and lower sums for increasing functions

a. Suppose the graph of a continuous function $f(x)$ rises steadily as x moves from left to right across an interval $[a, b]$. Let P be a partition of $[a, b]$ into n subintervals of length $\Delta x = (b - a)/n$. Show by referring to the accompanying figure that the difference between the upper and lower sums for f on this partition can be represented graphically as the area of a rectangle R whose dimensions are $[f(b) - f(a)]$ by Δx . (Hint: The difference $U - L$ is the sum of areas of rectangles whose diagonals $Q_0Q_1, Q_1Q_2, \dots, Q_{n-1}Q_n$ lie along the curve. There is no overlapping when these rectangles are shifted horizontally onto R .)

b. Suppose that instead of being equal, the lengths Δx_k of the subintervals of the partition of $[a, b]$ vary in size. Show that

$$U - L \leq |f(b) - f(a)| \Delta x_{\max},$$

where Δx_{\max} is the norm of P , and hence that $\lim_{\|P\| \rightarrow 0} (U - L) = 0$.

